ournals: Writing cmje.org/2006\_ur

International Coship credit should on and design, aca; 2) drafting the nal content; and rs should meet these criteria

s the Uniform and Figures. Leng 5000 words of body Case Report should deal cases. The case report should Abstract, Introduction, Case re-Figures. Abstract should be unstr ceed 150 words. These are limited 4 images. Editorial is usually written by Ed



## **BLOOD RESEARCH**

Volume 52 · Number 4 · December 2017 https://doi.org/10.5045/br.2017.52.4.235



## IL-1B-31 and IL-1Ra polymorphisms associated with increased host susceptibility to immune thrombocytopenia

Juwon Kim, M.D., Ph.D.

Department of Laboratory Medicine, Yonsei University Wonju College of Medicine, Wonju, Korea

Immune thrombocytopenia (ITP) is an autoimmune bleeding disorder in which the platelet autoantigens activate the immune system of the patient causing immune-mediated platelet destruction and/or suppression of platelet production. ITP can be either primary or secondary to other disorders, but the primary cause of the disease is still not well known. Meanwhile, many reports on genetic factors of ITP described several single nucleotide polymorphisms (SNPs), mostly inflammatory cytokine polymorphisms, associated with increased risk of ITP [1]. These SNPs may cause disturbance in the Th1 and Th2 cell balance leading to overproduction of inflammatory cytokines [2]. The SNPs in TNF- $\alpha$  gene are associated with increased phagocytic activity of macrophages and T-cytotoxic cells leading to the destruction of platelets via activation of apoptotic pathways [3]. Furthermore, TNF- $\alpha$  and - $\beta$  are associated with increased serum level of cytokines which plays a crucial role in the regulation of immune systems. Of note, the patient with  $TNF-\beta$  +252GG genotype was shown to be associated with platelet recovery in ITP patients after the eradication of Helicobacter pylori [4]. Moreover, the IFN-γ +874TT was found as a genotype that is associated with increased cytokine production which in turn causes increment of antibody production leading to increased platelet destruction [5].

In this issue of the **Blood Research**, Yadav et al. [6] evaluated the association of polymorphisms in interleukin (IL)-1B-31, IL-1B-511, and IL-1Ra with ITP, and revealed

that IL-1B-31 and IL-1Ra were significantly associated with ITP whereas IL-1B-511 failed to show association with ITP. Notably, a significant association of homozygous variant genotypes of IL-1B-31 was observed with severe ITP when compared with healthy controls and showed significant risk association with severe ITP, whereas IL-1Ra failed to show significant difference. Three polymorphisms in the promoter region of IL-1B, i.e. IL-1B-1464 (G/C; rs1143623; previously known as -1476), IL-1B-511 (C/T; rs16944), and IL-1B-31 (T/C; rs1143627), have been widely investigated, especially in association with gastric cancer risk and H. pylori infection. The IL-1 $\beta$  is a potent pro-inflammatory cytokine involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis [7]. Moreover, IL-1β is important in amplifying immune response to infection, and is a potent inhibitor of gastric acid secretion [8]. Although association studies of IL-1B polymorphism and autoimmune or immune-inflammatory diseases have been carried out, only few reports in relation to ITP are found. Wu et al. [9] described IL-1Ra but not IL-1B exon 5 polymorphism was associated with childhood ITP. Lack of the IL-1B-511T allele was associated with H. pylori infection in patients with early-onset ITP but not in those with late-onset ITP [10]. Most of the previously reported cases of IL-1B polymorphism in ITP were in relation to *H. pylori* infection. In the report by Yadav et al., in order to rule out ITP associated with secondary causes, patients with human immune deficiency virus, systemic lupus erythematous, and

236 Editorial

H. pylori were excluded from the study. Thus, direct association between ITP and IL-1B polymorphisms could be elucidated. Although the regulation of cytokine production is known to play a major role in the development of ITP, not many studies have focused on the host genetic factor as a potential triggering factor of the disease. The possible role of the IL-1B genotype in the development of ITP could be applied to determine the underlying etiology. However, the risk allele of IL-1B should be confirmed in other ethnic groups and should also be validated in larger cohort of patients before applying the concept since some instances the allele flips are observed in IL-1B depending on the pathogenesis of disease.

## **REFERENCES**

- Rezaeeyan H, Jaseb K, Alghasi A, Asnafi AA, Saki N. Association between gene polymorphisms and clinical features in idiopathic thrombocytopenic purpura patients. Blood Coagul Fibrinolysis 2017;28:617-22.
- 2. Morgan DS, Afifi RA, El-Hoseiny SM, Amin DG, Ibrahim SYG. The potential association of tumor necrosis factor- $\beta$ eta (252 G/A) cytokine gene polymorphism with immune thrombocytopenic purpura among Egyptian children. Hematology 2017:1-5.
- 3. El Sissy MH, El Sissy AH, Elanwary S. Tumor necrosis factor- $\alpha$ -308G/A gene polymorphism in Egyptian children with immune

- thrombocytopenic purpura. Blood Coagul Fibrinolysis 2014;25: 458-63
- Suzuki T, Matsushima M, Shirakura K, et al. Association of inflammatory cytokine gene polymorphisms with platelet recovery in idiopathic thrombocytopenic purpura patients after the eradication of Helicobacter pylori. Digestion 2008;77:73-8.
- 5. Lee YH, Bae SC. Association between interferon-γ+874 T/A polymorphism and susceptibility to autoimmune diseases: a meta-analysis. Lupus 2016;25:710-8.
- 6. Yadav DK, Tripathi AK, Gupta D, et al. Interleukin-1B (*IL-1B-31* and *IL-1B-511*) and interleukin-1 receptor antagonist (*IL-1Ra*) gene polymorphisms in primary immune thrombocytopenia. Blood Res 2017;52:264-9.
- Dinarello CA. Biologic basis for interleukin-1 in disease. Blood 1996;87:2095-147.
- 8. Schneider BG, Camargo MC, Ryckman KK, et al. Cytokine polymorphisms and gastric cancer risk: an evolving view. Cancer Biol Ther 2008;7:157-62.
- Wu KH, Peng CT, Li TC, Wan L, Tsai CH, Tsai FJ. Interleukin-1beta exon 5 and interleukin-1 receptor antagonist in children with immune thrombocytopenic purpura. J Pediatr Hematol Oncol 2007;29:305-8.
- Satoh T, Pandey JP, Okazaki Y, et al. Single nucleotide polymorphism of interleukin-1beta associated with Helicobacter pylori infection in immune thrombocytopenic purpura. Tissue Antigens 2009;73:353-7.